

Reduce fibrosis and adhesions following tendon and peripheral nerve surgery

Protect Your Patient



Consider the importance to your patient of reducing adhesions.

Separate and protect tissues

Designed to coat, separate and protect tissues where postoperative adhesions are likely to form.

Reduce fibrosis and adhesions

Studies have shown that coating tissues with a gel barrier of CMC/PEO can reduce the formation of adhesions.

Indications

- Tendon & Nerve Repair
- Tenolysis & Neurolysis
- Brachial Plexus Reconstruction
- MCP & PIP Joint Arthroplasty
- Trigger Finger & De Quervain Syndromes
- Dupuytren's Contracture
- Carpal Tunnel Release

Long history of safe and effective tissue protection.*

DYNAVISC® is a sterile, absorbable combination of carboxymethylcellulose (CMC) and polyethylene oxide (PEO).

2 Year Shelf Life

Introducing
DYNAVISC®

Adhesion Barrier Gel



*Based on safety and effectiveness experience of FzioMed, Inc. gel in spinal surgery (over 250,000 surgeries), and independent clinical and pre-clinical studies. Rhyne A, et al. Spine, 37 (8):631-641, 2012. Franssen P. Ann Sur Innov Res, 2:2, 2008. Assietti R, et al. Spine, 33 (16): 1762-1765, 2008. Arnold P, et al. AANS 2008. Guizzardi G, et al. Congress of Neurological Surgeons (CNS) 2006. Rodgers, KE, et al. Spine J, 3:277-284, 2003. Rodgers KE, et al. Congress of Neurological Surgeons (CNS) 2003. Data on file, FzioMed, Inc.

FzioMed® and Dynavisc® are registered trademarks of FzioMed, Inc.
MM048(-)

fzioMed

www.fziomed.com